

ON ASYMPTOTIC EXTENSION DIMENSION

DUŠAN REPOVŠ AND MYKHAILO ZARICHNYI

ABSTRACT. The aim of this paper is to introduce an asymptotic counterpart of the extension dimension defined by Dranishnikov. The main result establishes a relation between the asymptotic extensional dimension of a proper metric space and extension dimension of its Higson corona.

1. INTRODUCTION

Asymptotic dimension defined by Gromov [7] has been an object of study in numerous publications (see expository paper [2]). A metric space (X, d) is of asymptotic dimension $\leq n$ (written $\text{asdim } X \leq n$) if for every $D > 0$ there exists a uniformly bounded cover \mathcal{U} of X such that $\mathcal{U} = \mathcal{U}^0 \cup \dots \cup \mathcal{U}^n$, where every family \mathcal{U}^i is D -disjoint, $i = 0, 1, \dots, n$. Recall that a family \mathcal{A} of subsets of X is *uniformly bounded* if

$$\text{mesh } \mathcal{A} = \sup\{\text{diam } A \mid A \in \mathcal{A}\} < \infty$$

and is called *D-disjoint* if $\inf\{d(a, a') \mid a \in A, a' \in A'\} > D$, for every distinct $A, A' \in \mathcal{A}$.

The asymptotic dimension can be characterized in different terms; in particular, in terms of extension of maps into euclidean spaces. The aim of this paper is to introduce an asymptotic analogue of the extension dimension introduced by Dranishnikov in [3, 4].

2. PRELIMINARIES

A typical metric is denoted by d . By $N_r(x)$ we denote the open ball of radius r centered at a point x of a metric space.

2.1. Asymptotic category. A map $f: X \rightarrow Y$ between metric spaces is called (λ, ε) -Lipschitz for $\lambda > 0$, $\varepsilon \geq 0$ if $d(f(x), f(x')) \leq \lambda d(x, x') + \varepsilon$ for every $x, x' \in X$. A map is called *asymptotically Lipschitz* if it is (λ, ε) -Lipschitz for some $\lambda, \varepsilon > 0$. The $(\lambda, 0)$ -Lipschitz maps are also called λ -Lipschitz, $(1, 0)$ -Lipschitz maps are also called *short*. A metric space X is called *proper* if every closed ball in X is compact.

The *asymptotic category* \mathcal{A} was introduced by Dranishnikov [4]. The objects of \mathcal{A} are proper metric spaces and the morphisms are proper asymptotically Lipschitz maps. Recall that a map is called *proper* if the preimage of every compact set is compact.

We also need the notion of a *coarse map*. A map between proper metric spaces is called *coarse uniform* if for every $C > 0$ there is $K > 0$ such that for every $x, x' \in X$ with $d(x, x') < C$ we have $d(f(x), f(x')) < K$. A map $f: X \rightarrow Y$ is called *metric proper* if the preimage $f^{-1}(B)$

Date: January 26, 2013.

2010 *Mathematics Subject Classification.* Primary: 54F45, 54C40; Secondary: 55M10.

Key words and phrases. Asymptotic category, asymptotic extension dimension, Higson compactification, Higson corona, coarse uniform map, coarse CW complex, proper asymptotically Lipschitz map, slowly oscillating map, almost geodesic space, Lipschitz neighborhood euclidean extensor, foliated warped cone.

This research was supported by the Slovenian Research Agency grants P1-0292-0101 and J1-2057-0101.

is bounded for every bounded set $B \subset Y$. A map is *coarse* if it is metric proper and coarse uniform.

2.2. Higson compactification and Higson corona. Let $\varphi: X \rightarrow \mathbb{R}$ be a function defined on a metric space X . For every $x \in X$ and every $r > 0$ let $V_r\varphi(x) = \sup\{|\varphi(y) - \varphi(x)| \mid y \in N_r(x)\}$. A function φ is called *slowly oscillating* whenever for every $r > 0$ we have $V_r\varphi(x) \rightarrow 0$ as $x \rightarrow \infty$ (the latter means that for every $\varepsilon > 0$ there exists a compact subspace $K \subset X$ such that $|V_r\varphi(x)| < \varepsilon$ for all $x \in X \setminus K$).

Let \bar{X} be the compactification of X that corresponds to the family of all continuous bounded slowly oscillation functions. The *Higson corona* of X is the remainder $\nu X = \bar{X} \setminus X$ of this compactification. It is well-known that the Higson corona is a functor from the category of proper metric space and coarse maps into the category of compact Hausdorff spaces. In particular, if $X \subset Y$, then $\nu X \subset \nu Y$.

For any subset A of X we denote by A' its trace on νX , i. e. the intersection of the closure of A in \bar{X} with νX . Obviously, the set A' coincides with the Higson corona νA .

2.3. Cones. Let X be a metric space of diameter ≤ 1 . The *open cone* of X is the set $\mathcal{O}X = (X \times \mathbb{R}_+)/ (X \times \{0\})$ endowed with the metric (by $[x, t]$ we denote the equivalence class of $(x, t) \in X \times \mathbb{R}_+$):

$$d([x_1, t_1], [x_2, t_2]) = |t_1 - t_2| + \min\{t_1, t_2\}d(x_1, x_2).$$

For a map $f: X \rightarrow Y$ of metric spaces we denote by $\mathcal{O}f: \mathcal{O}X \rightarrow \mathcal{O}Y$ the map defined as $\mathcal{O}f([x, t]) = [f(x), t]$.

Proposition 2.1. *If $f: X \rightarrow Y$ is a Lipschitz map then $\mathcal{O}f$ is an asymptotically Lipschitz map.*

Proof. Suppose a map $f: X \rightarrow Y$ is λ -Lipschitz. Then for any $[x_1, t_1], [x_2, t_2] \in \mathcal{O}X$ we have

$$\begin{aligned} d(\mathcal{O}f([x_1, t_1]), \mathcal{O}f([x_2, t_2])) &= d([f(x_1), t_1], [f(x_2), t_2]) \\ &= |t_1 - t_2| + \min\{t_1, t_2\}d(f(x_1), f(x_2)) \\ &\leq \lambda'(|t_1 - t_2| + \min\{t_1, t_2\}d(x_1, x_2)), \end{aligned}$$

where $\lambda' = \max\{\lambda, 1\}$. □

The open cone of a finite CW-complex is a coarse CW-complex in the sense of [8].

Denote by $\alpha_L: \mathcal{O}L \rightarrow \mathbb{R}$ the function defined by $\alpha_L([x, t]) = t$. Obviously, α_L is a short function. Let $\tilde{\mathcal{O}}L = \{[x, t] \in \mathcal{O}L \mid t \geq 1\}$. Denote by $\beta_L: \tilde{\mathcal{O}}L \rightarrow L$ the map $\beta_L([x, t]) = x$.

Lemma 2.2. *The map β_L is slowly oscillating.*

Proof. For $R > 0$, the R -ball centered at $[x, 0]$ is $\{[x, t] \mid t < R\}$. If $d([x, t], [x_1, t_1]) < K < R$, then $|t - t_1| + \min\{t, t_1\}d(x, x_1) < K$, i.e. $(t - R)d(x, x_1) < R$ and $d(x, x_1) < K/(t - K)$. Therefore, $d(\beta_L(x), \beta_L(x_1)) < K/(R - K) \rightarrow 0$ as $R \rightarrow \infty$. □

Let $\bar{\beta}_L: \tilde{\mathcal{O}}L \rightarrow L$ be the (unique) extension of the map β_L . Denote by $\eta_L: \nu\tilde{\mathcal{O}}L \rightarrow L$ the restriction of $\bar{\beta}_L$.

Proposition 2.3. *Let $f: A \rightarrow \mathcal{O}L$ be a proper asymptotically Lipschitz map defined on a proper closed subset A of a proper metric space X . There exists a neighborhood W of A in X , a proper asymptotically Lipschitz map $g: W \rightarrow \mathcal{O}L$ with the following property: there exist constants $\lambda, s > 0$ such that $\alpha_L(g(a)) \leq \lambda d(a, X \setminus W) + s$.*

Proof. We may assume that L is a subset of I^n for some n and there exists a Lipschitz retraction $r: U \rightarrow L$ of a neighborhood U of L in I^n . Since $\mathcal{O}I^n$ is Lipschitz equivalent to \mathbb{R}_+^{n+1} , there exists a (λ', s') -Lipschitz extension $\tilde{g}: X \rightarrow \mathcal{O}I^n$ of g .

Put $W = \tilde{g}^{-1}(\mathcal{O}U)$ and $\bar{g} = \tilde{g}|_W$. For every $a \in A$ and $w \in X \setminus W$ we have

$$d(g(a), \tilde{g}(w)) \leq \lambda' d(a, w) + s' \leq \lambda' d(a, X \setminus W) + s.$$

Suppose that $d(L, I^n \setminus U) = c > 0$, then, since $\tilde{g}(w) \notin CU$,

$$\begin{aligned} d(g(a), \tilde{g}(w)) &= |\alpha_L(g(a)) - \alpha_L(\tilde{g}(w))| \\ &\quad + \min\{\alpha_L(g(a)), \alpha_L(\tilde{g}(w))\} d(\beta_L(g(a)), \beta_L(\tilde{g}(w))) \\ &\geq |\alpha_L(g(a)) - \alpha_L(\tilde{g}(w))| + c \min\{\alpha_L(g(a)), \alpha_L(\tilde{g}(w))\} \\ &\geq c' \alpha_L(g(a)), \end{aligned}$$

where $c' = \min\{c, 1\}$. Then $\alpha_L(g(a)) \leq \lambda d(a, X \setminus W) + s$, where $\lambda = \lambda'/c'$, $s = s'/c'$. \square

3. AUXILIARY RESULTS

In this section we shall collect some results needed for the proof of the main result. They are proved in [4] but it turns out that we have also to cover the case of functions with infinite values.

A map $f: X \rightarrow \mathbb{R}_+ \cup \{\infty\}$ is said to be *coarsely proper* if the preimage $f^{-1}([0, c])$ is bounded for every $c \in \mathbb{R}_+$.

Lemma 3.1. *For any function $\varphi: X \rightarrow \mathbb{R}_+$ with $\varphi(x) \rightarrow 0$ as $x \rightarrow \infty$ the function $1/\varphi: X \rightarrow \mathbb{R}_+ \cup \{\infty\}$ is coarsely proper.*

Proposition 3.2. *Let $f: X \rightarrow \mathbb{R}_+ \cup \{\infty\}$ be a coarsely proper function. There exists an asymptotically Lipschitz proper function $q: X \rightarrow \mathbb{R}_+$ with $q \leq f$.*

Proof. This was proved in [4] for the case of $f: X \rightarrow \mathbb{R}_+$ (see Proposition 3.5). That proof also works in our case. \square

Proposition 3.3. *Let $f_n: X \rightarrow \mathbb{R}_+ \cup \{\infty\}$ be a sequence of coarsely proper functions. Then there exists a filtration $X = \cup_{n=1}^\infty A_n$ and a coarsely proper function $f: X \rightarrow \mathbb{R}_+$ with $f|_{A_n} \leq n$ and $f|(X \setminus A_n) \leq f_n$ for every n .*

Proof. Let $B_n = \cup_{i=1}^n f_i^{-1}([0, n])$. the sets B_i are bounded and $B_1 \subset B_2 \subset \dots$. Therefore, there exist bounded subsets $A_1 \subset A_2 \subset \dots$ such that $A_n \cap (\cup_{i=1}^\infty B_i) = B_n$ and $\cup_{i=1}^\infty A_i = X$. For $x \in A_n \setminus A_{n-1}$, put $f(x) = n$. Obviously, f is coarsely proper and $f|_{A_n} \leq n$. Now suppose that $x \notin A_n$, then $x \notin B_n$ and therefore $x \notin f_n^{-1}([0, n])$, i.e. $f_n(x) > n \geq f|(X \setminus A_n)$. \square

The following is an easy modification of Lemma 3.6 from [4] and the proof of it works in our case as well.

Lemma 3.4. *Suppose that $f: A \rightarrow \mathbb{R}_+ \cup \{\infty\}$ is a coarsely proper map defined on a closed subset A of a proper metric space X and $g: W \rightarrow \mathbb{R}_+$ is a proper asymptotically Lipschitz map such that $g \leq f|_W$ and there exist λ, s such that $\lambda d(a, X \setminus W) + s \geq g(a)$ for every $a \in A$. Then there exists a proper asymptotically Lipschitz map $\bar{g}: X \rightarrow \mathbb{R}_+$ for which $\bar{g} \leq f$ and $\bar{g}|_A = g$.*

3.1. Almost geodesic spaces. A metric space X is said to be *almost geodesic* if there exists $C > 0$ such that for every two points $x, y \in X$ there is a short map $f: [0, Cd(x, y)] \rightarrow X$ with $f(0) = x$, $f(Cd(x, y)) = y$. If in this definition $C = 1$, then we come to the well-known notion of *geodesic* space.

We are going to describe a construction of embedding of a discrete metric space X into an almost geodesic space of the asymptotic dimension $\min\{\text{asdim} X, 1\}$. For an unbounded discrete metric space X with base point x_0 define a function $f: X \rightarrow [0, \infty)$ by the formula $f(x) = d(x, x_0)$. Choose a sequence $0 = t_0 < t_1 < t_2 < \dots$ in $f(X)$ so that $t_{i+1} > 2t_i$ for every i . To every pair of points $x, y \in f^{-1}([t_i, t_{i+1}])$, for some i , attach the line segment $[0, d(x, y)]$ along its endpoints. Let \hat{X} is the union of X and all attached segments. We endow \hat{X} with the maximal metric that agrees with the initial metric on X and the standard metric on every attached segment.

Note that since X is discrete and proper, every set $f^{-1}([t_i, t_{i+1}])$ is finite and therefore \hat{X} is a proper metric space.

Proposition 3.5. *The space \hat{X} is almost geodesic.*

Proof. Suppose that $x, y \in \hat{X}$, then $x \in [x_1, x_2]$, $y \in [y_1, y_2]$, where $x_1, x_2, y_1, y_2 \in X$ and $[x_1, x_2]$, $[y_1, y_2]$ are attached segments. We may suppose that $d(x, y) = d(x, x_1) + d(x_1, y_1) + d(y_1, y)$.

Case 1): There exists i such that $x_1, y_1 \in f^{-1}([t_i, t_{i+1}])$. Then $[x, x_1] \cup [x_1, y_1] \cup [y_1, y]$ is a segment of diameter $d(x, y)$ that connects x and y in \hat{X} .

Case 2): $f(x_1) \in [t_i, t_{i+1}]$, $f(y_1) \in [t_j, t_{j+1}]$, where $i \neq j$. Without loss of generality, we may assume that $i < j$.

Obviously, $d(x_1, y_1) \leq d(x, y)$. Since $|t_j - t_{j-1}| \leq d(x_1, y_1)$, we see that $|t_j - t_{j-1}| \leq d(x, y)$. This implies that $t_j/2 \leq d(x, y)$, or equivalently, $t_j \leq d(x, y)$. Besides, $d(y_1, f^{-1}([0, t_{j-1}])) \leq d(x_1, y_1) \leq d(a, b)$.

For every $k = i, i+1, \dots, j_1$ choose $z_k \in f^{-1}(t_k)$. Then

$$\begin{aligned} d(y_1, z_{j-1}) &\leq d(y_1, f^{-1}([0, t_{j-1}])) + \text{diam}(f^{-1}([0, t_{j-1}])) \\ &\leq d(a, b) + 2t_{j-1} \leq d(a, b) + t_j \leq 3d(a, b). \end{aligned}$$

We connect x and y by the segment

$$J = [x, x_1] \cup [x_1, z_1] \cup \bigcup_{k=i}^{j-1} [z_k, z_{k+1}] \cup [z_{j-1}, y_1] \cup [y_1, y].$$

Then

$$\begin{aligned} \text{diam } J &\leq d(x, x_1) + d(x_1, z_{i+1}) + \left(\sum_{k=i+1}^{j-1} d(z_k, z_{k+1}) \right) + d(z_{j-1}, y_1) + d(y_1, y) \\ &= d(x, y) + 2t_{i+1} + \sum_{k=i+1}^{j-1} 2t_{k+1} + 5d(x, y) + d(x, y) \\ &\leq 7d(x, y) + 2(t_{i+1} + \dots + t_j) \leq 7d(x, y) + 4t_j \\ &\leq 15d(x, y). \end{aligned}$$

□

We need a version of the fact proved in [4] for geodesic spaces.

Proposition 3.6. *Let $f: X \rightarrow Y$ be a coarse uniform map of an almost geodesic space X . Then f is asymptotically Lipschitz.*

Proof. Let C be a constant from the definition of almost geodesic space. Suppose $x, y \in X$, then there exists a short map $\alpha: [0, Cd(x, y)] \rightarrow X$ such that $\alpha(0) = x$, $\alpha(Cd(x, y)) = y$. There exist points $0 = t_0 < t_1 < \dots < t_{k-1} < t_k = Cd(x, y)$, where $k \leq [d(x, y)] + 1$, such that $|t_i - t_{i-1}| \leq C$ for every $i = 1, \dots, k$.

Since f is coarse uniform, there exists $R > 0$ such that $d(f(x'), f(y')) < R$ whenever $d(x', y') \leq C$. Then

$$\begin{aligned} d(f(x), f(y)) &\leq \sum_{i=1}^k d(f(\alpha(t_i)), f(\alpha(t_{i-1}))) \leq kR \leq ([d(x, y)] + 1)R \\ &\leq Rd(x, y) + 2R. \end{aligned}$$

□

4. ASYMPTOTIC EXTENSION DIMENSION

Let P be an object of the category \mathcal{A} . For any object X of \mathcal{A} the *Kuratowski notation* $X\tau P$ means the following: for every proper asymptotically Lipschitz map $f: A \rightarrow P$ defined on a closed subset A of X there is a proper asymptotically Lipschitz extension of f onto X .

Denote by \mathcal{L} the class of compact absolute Lipschitz neighborhood euclidean extensors (ALNER). Following [], we define a preorder relation \leq on \mathcal{L} . For $L_1, L_2 \in \mathcal{L}$, we have $L_1 \leq L_2$ if and only if $X\tau\mathcal{O}L_1$ implies $X\tau\mathcal{O}L_2$ for all proper metric spaces X . This preorder relation leads to the following equivalence relation \sim on \mathcal{L} : $L_1 \sim L_2$ if and only if $L_1 \leq L_2$ and $L_2 \leq L_1$. We denote by $[L]$ the equivalence class containing $L \in \mathcal{L}$.

For a proper metric space X , we say that its *asymptotic extension dimension does not exceed* $[\mathcal{O}L]$ (briefly $\text{as-ext-dim}X \leq [\mathcal{O}L]$) whenever $X\tau\mathcal{O}L$. If $\text{as-ext-dim}X \leq [\mathcal{O}L]$, then the equality $\text{as-ext-dim}X = [\mathcal{O}L]$ means the following. If we also have $\text{as-ext-dim}X \leq [\mathcal{O}L']$, then $[\mathcal{O}L] \leq [\mathcal{O}L']$. By [4] (see also [11]), the element $[*]$ is maximal.

Theorem 4.1. *Let L be a compact metric ALNER. The following conditions are equivalent:*

- (1) $\text{as-ext-dim}X \leq [\mathcal{O}L]$;
- (2) $\text{ext-dim}\nu X \leq [L]$.

Proof. Assume that $\text{as-ext-dim}X \leq [\mathcal{O}L]$. Let $\varphi: C \rightarrow L$ be a map defined on a closed subset C of νX . Since $L \in \text{ANE}$, there exists an extension $\varphi': V \rightarrow L$ of φ over a closed neighborhood V of C in $\bar{X} = X \cup \nu X$. Then $\text{Var}_R \varphi'(x) \rightarrow 0$ as $x \rightarrow \infty$, for any fixed $R > 0$. By Lemma 3.1, the function

$$f_n: V \cap X \rightarrow \mathbb{R}_+ \cup \{\infty\}, \quad f_n(x) = \frac{1}{\text{Var}_R \varphi'(x)},$$

is coarsely proper, for every $n \in \mathbb{N}$. By Proposition 3.3, there is a coarsely proper function $f: V \cap X \rightarrow \mathbb{R}_+$ and a filtration $V \cap X = \bigcup_{n=1}^{\infty} A_n$ such that $f|_{A_n} \leq n$ and $f|(X \setminus A_n) \leq f_n$. By Proposition 3.5 from [4], there is an asymptotically Lipschitz function $q: V \cap X \rightarrow \mathbb{R}_+$ with $q \leq f$. We suppose that q is (λ, s) -Lipschitz for some $\lambda, s > 0$. Define the map $g: V \cap X \rightarrow \mathcal{O}L$ by the formula $g(x) = [\varphi'(x), q(x)]$.

We are going to check that the map $g(x)$ is asymptotically Lipschitz. Let $x, y \in V \cap X$ and $n - 1 \leq d(x, y) \leq n$.

Suppose that $x, y \in (V \cap X) \setminus A_n$, then $q(x) \leq f_n(x)$, $q(y) \leq f_n(y)$. We have

$$\begin{aligned} d(g(x), g(y)) &= |q(x) - q(y)| + \min\{q(x), q(y)\} d(\varphi'(x), \varphi'(y)) \\ &\leq \lambda d(x, y) + s + \min\{q(x), q(y)\} \text{Var}_n \varphi'(x) \\ &\leq \lambda d(x, y) + s + 1. \end{aligned}$$

If $x \in A_n$, then $q(x) \leq n$ and we obtain

$$\begin{aligned} d(g(x), g(y)) &\leq \lambda d(x, y) + s + nd(\varphi'(x), \varphi'(y)) \\ &\leq \lambda d(x, y) + s + nd \operatorname{diam} L \leq \lambda d(x, y) + s + (d(x, y) + 1) \operatorname{diam} L \\ &\leq (\lambda + \operatorname{diam} L) d(x, y) + (s + \operatorname{diam} L). \end{aligned}$$

We argue similarly if $y \in A_n$. Now, by the assumption, there is an asymptotically Lipschitz extension $\bar{g}: X \rightarrow \mathcal{OL}$ of g . Consider the composition $\eta_L \nu \bar{g}: \nu X \rightarrow \mathcal{OL}$. Obviously, $\eta_L \nu \bar{g}|C = \varphi$.

Let $f: A \rightarrow \mathcal{OL}$ be an asymptotically Lipschitz map defined on a proper closed subset A of a proper metric space X . By Proposition 2.2, there is a proper asymptotically Lipschitz map $\tilde{f}: W \rightarrow \mathcal{OL}$ and constants λ, s such that $\alpha_L f(a) \leq \lambda d(a, X \setminus W) + s$ for all $a \in A$. Denote by $\varphi: \nu X \rightarrow L$ an extension of the composition $\eta_L \nu \tilde{f}$. Since L is an absolute neighborhood extensor, there exists an extension $\psi: V \rightarrow L$ of φ onto a closed neighborhood of νX in the Higson compactification \bar{X} . Extend ψ to a map $\hat{\psi}: (V \cap X)^\circ \rightarrow L$ as follows. Let J be a segment attached to V with endpoints a and b . We require that $\hat{\psi}$ linearly maps J onto a geodesic segment in L with endpoints $\psi(a)$ and $\psi(b)$.

We show that $\hat{\psi}$ is a slowly oscillating map. Since ψ is slowly oscillating, for every $\varepsilon > 0$ and $R > 0$ there exists $K > 0$ such that $\operatorname{Var}_R \psi(x) < \varepsilon$ whenever $d(x, x_0) > K$. Suppose that $\hat{\psi}$ is not slowly oscillating, then there exist $R > 0$, $C > 0$, and sequences (x_1^i) , (x_2^i) in $(V \cap X)^\circ$ such that $d(x_1^i, x_2^i) < R$, $x_1^i \rightarrow \infty$, $x_2^i \rightarrow \infty$ and $d(\hat{\psi}(x_1^i), \hat{\psi}(x_2^i)) > C$ for every i . We assume that $x_1^i \in [a_1^i, b_1^i]$, $x_2^i \in [a_2^i, b_2^i]$, for every i , where $a_1^i, b_1^i, a_2^i, b_2^i \in X \cap V$. Without loss of generality we may assume that $a_1^i \rightarrow \infty$ and there exists $C_1 > 0$ such that $d(\hat{\psi}(x_1^i), \hat{\psi}(a_1^i)) > C_1$ for every i . If $d(a_1^i, b_1^i) < K$ for all i and some $K > 0$, then $d(\hat{\psi}(x_1^i), \hat{\psi}(a_1^i)) < d(\hat{\psi}(a_1^i), \hat{\psi}(b_1^i)) \rightarrow 0$, and we obtain a contradiction. Therefore, we may assume that $d(a_1^i, b_1^i) \rightarrow \infty$. Then $d(a_1^i, x_1^i)/d(a_1^i, b_1^i) < R/d(a_1^i, b_1^i) \rightarrow 0$ and therefore, by the definition of the map $\hat{\psi}$, $d(\hat{\psi}(x_1^i), \hat{\psi}(a_1^i))/d(\hat{\psi}(a_1^i), \hat{\psi}(b_1^i)) \rightarrow 0$. Then obviously $d(\hat{\psi}(x_1^i), \hat{\psi}(a_1^i)) \rightarrow 0$ and we obtain a contradiction.

Since the map \tilde{f} is asymptotically Lipschitz, there exists $K > 0$ such that for any $a \in W$ we have

$$\operatorname{diam}(\alpha_L \tilde{f}(N_1(a)) + \alpha_L \tilde{f}(a) \operatorname{diam}(\psi(N_1(a))) \leq K.$$

Define the function $r: (X \cap V)^\circ \rightarrow \mathbb{R}_+ \cup \{\infty\}$ by the formula $r(x) = K/(\psi(N_1(x)))$. We have $f(a) \leq r(a)$ for every $a \in A$. The function r is asymptotically proper and by Proposition 3.2, there exists a (λ', s') -Lipschitz function $\bar{f}: X \rightarrow \mathbb{R}_+$, for some λ', s' , with $\bar{f} \leq r$ and $\bar{f}|A = \alpha_L f$.

Define a map $g: (X \cap V)^\circ \rightarrow \mathbb{R}$ by the formula $g(x) = (\psi(x), \bar{f}(x))$. Obviously, $g|A = f$. We are going to show that g is a coarse uniform map. Suppose $x, y \in X$, $d(x, y) < 1$, then

$$\begin{aligned} d(g(x), g(y)) &\leq |\bar{f}(x) - \bar{f}(y)| + \min\{\bar{f}(x), \bar{f}(y)\} d(\psi(x), \psi(y)) \\ &\leq \lambda' + s' + K. \end{aligned}$$

Note that, since \bar{f} is proper, g is also proper. Since g is coarse uniform, by Proposition 3.6, g is asymptotically Lipschitz. \square

Corollary 4.2. (Finite Sum Theorem) *Suppose X is a proper metric space, $X = X_1 \cup X_2$, where X_1, X_2 are closed subsets of X with $\operatorname{as-ext-dim} X_i \leq [\mathcal{OL}]$, $i = 1, 2$, for some $L \in \mathcal{L}$. Then $\operatorname{as-ext-dim} X \leq [\mathcal{OL}]$.*

Proof. Since $\nu X = \nu X_1 \cup \nu X_2$, the result follows from Theorem 4.1 and the finite sum theorem for extension dimension (see [2]). \square

5. REMARKS AND OPEN QUESTIONS

Question 5.1. Does the equality $\text{as-ext-dim}\mathbb{R}^n = S^n$ hold?

Question 5.2. Let L_1, L_2 be finite polyhedra in euclidean spaces endowed with the induced metric. Is the inequality $[L_1] \leq [L_2]$ introduced in [3] equivalent to the inequality $[L_1] \leq [L_2]$ as in Section 4?

One can define analogue of the asymptotic extension dimension by using warped cones instead of open cones. Following [9] we review this construction briefly. Let \mathcal{F} be a foliation on a compact smooth manifold V . Let N be any complementary subbundle to $T\mathcal{F}$ in TM . Choose Euclidean metrics g_N in N and $g_{\mathcal{F}}$ in $T\mathcal{F}$. The *foliated warped cone* $\mathcal{O}_{\mathcal{F}}$ is the manifold $V \times [0, \infty)/V \times \{0\}$ equipped with the metric induced for $t \geq 1$ by the Riemannian metric $g_R + g_{\mathcal{F}} + t^2g_N$. The metric structure on any bounded neighborhood of the distinguished point is irrelevant.

Question 5.3. Is the obtained warped cone an absolute neighborhood extensor in the asymptotic category?

An affirmative answer to this question would allow us to define asymptotic extension dimension theory with the values in warped cones.

REFERENCES

- [1] G. Bell and A. N. Dranishnikov, *On asymptotic dimension of groups*, Algebraic and Geometric Topology **1** (2001), 57–71.
- [2] G. Bell and A. N. Dranishnikov, *Asymptotic dimension*, preprint on the Archives <http://arxiv.org/abs/math/0703766v2>
- [3] A. N. Dranishnikov, *On the theory of extensions of mappings of compact spaces*, (Russian) Uspekhi Mat. Nauk **53**:5 (1998), 65–72; English transl. in Russian Math. Surveys **53**:5 (1998), 929–935.
- [4] A. N. Dranishnikov, *Asymptotic topology*, Russian Math. Surveys **55**(2000), No 6, 71–116.
- [5] A. Dranishnikov and J. Dydak, *Extension dimension and extension types*, Trudy Mat. Inst. Steklova **212** (1996), Oتبrazh. i Razmer., 61–94; English transl. in Proc. Steklov Inst. Math. **212**:1 (1996), 55–88.
- [6] A. Dranishnikov and J. Dydak, *Extension theory of separable metrizable spaces with applications to dimension theory*, Trans. Amer. Math. Soc. **353**:1 (2001) 133–156.
- [7] M. Gromov, *Asymptotic invariants of infinite groups*, Geometric group theory, Vol. 2 (Sussex, 1991), London Math. Soc. Lecture Note Ser., vol. 182, Cambridge Univ. Press, Cambridge, 1993, pp. 1–295.
- [8] P. Mitchener, *Coarse homology theories*, Algebr. Geom. Topol. **1** (2001), 271–297 (electronic).
- [9] J. Roe, *From foliations to coarse geometry and back*, in: Analysis and geometry in foliated manifolds, (X. Masa, E. Macias-Virgós, J.A. Alvarez López, Editors), World Scientific, Singapore 1995, pp. 195–206.
- [10] J. Roe, *Warped cones and property A*, Geometry & Topology **9** (2005) 163178 Published: 6 January 2005, Corrected: 7 March 2005
- [11] M. Sawicki, *Absolute extensors and absolute neighborhood extensors in asymptotic categories*, Topology Appl. **150**:13 (2005), 59–78.

FACULTY OF MATHEMATICS AND PHYSICS, AND FACULTY OF EDUCATION, UNIVERSITY OF LJUBLJANA, JADRANSKA 19, 1000 LJUBLJANA, SLOVENIA
E-mail address: `dusan.repovs@guest.arnes.si`

DEPARTMENT OF MECHANICS AND MATHEMATICS, LVIV NATIONAL UNIVERSITY, UNIVERSYTETSKA 1, 79000 LVIV, UKRAINE
E-mail address: `mzar@litech.lviv.ua`